Halo-Gravity Traction for the Treatment of Cervical Spine Disorders

Bram Verhofste MD, Michael Timothy Hresko MD, John Emans MD, Michael Glotzbecker MD, Brian Snyder MD PhD, Craig Birch MD, Nora O'Neill BA, Lawrence Karlin MD, Mark Proctor MD, Daniel Hedequist MD

13th International Congress on Early Onset Scoliosis November 20-22, 2019 – Atlanta, GA



Boston Children's Hospital Orthopedic Center







Disclosures

I have <u>no relevant financial relationships</u> with the manufacturer(s) of any commercial product(s) and/or provider(s) of commercial services



Boston Children's Hospital Orthopedic Center





Case Example

Issues

- Young age (size, compliance)
- Neurologic deficits
- Poor head control
- Significant kyphosis

1 yo M w/ NF1 + rhabdoid tumor

- Tumor resection + laminectomies
- CT + RT













Case

2 years later

Experimental CT Loss of head control + ambulation Failed bracing





Problems

Loss of posterior elements Small anatomy Radiated bone

- Poor quality
- Poor growth potential

Definitive solution ??



Case





Boston Children's Hospital Orthopedic Center





Objectives

- 1. HGT efficacy
- 2. <u>Safety + complications</u>
- 3. <u>Protocol</u> for pediatric cervical spines disorders





Boston Children's Hospital Orthopedic Center





Methods

Study Design: Retrospective cohort Inclusion

- Age <25 years

- Cervical spine deformity
- HGT



Cervical deformity

Cervical kyphosis Atlantoaxial rotatory subluxation Basilar invagination Os odontoideum Cervical arthritis causing deformity



<u>Data</u>

- Correctability
- Mod. Clavien-Dindo-Sink classification









Methods: HGT Protocol

** n=3 HVGT

Traction Details

- Halo-crown w/ 4-8 pins
- 1-2 lbs / day
- 20%-50% BW
- Pin-site care / neuro exams

Duration

- Variable
- 24/7 traction (excl. bathroom / shower)
- No activity restrictions













Results: Demographics

Cohort: 28 cases (mean 11.3 yo)





Results: HGT Details

Ou <u>tcom</u> e	es	
Characteristic	Mean	IQR
% Body weight (<i>kg</i>)	29%	19%-40%
HGT duration (<i>days</i>)	25	13-29
Length of stay (<i>day</i> s)	35	17-43
Surgical stabilization (<i>n; %</i>)	25	89%
	1	000
ě - 1	j ×.	g
		Je l
Pre	- T 3d	
Boston Children's Hospital		RVARD MEDICAL
Orthopedic Center	TEA	CHING HOSPITA

Results: HGT Efficacy



Methods: HGT Complications

9 AE (32%)

Grade I

- 4 pin-site infections (14%)
- 4 transient paresthesia's (14%)

Grade II

- 1 flaccid paralysis (4%)
- (T21 w/ Os, SCC, and myelopathy)

No major Grade III-V AE's



Boston Children's Hospital Orthopedic Center



HARVARD MEDICAL SCHOOL TEACHING HOSPITAL



Minimal clinical relevance

Weight removal + Reduced progression

Limitations



Conclusions

Efficacious + safe

Grade I AE's

Decreased surgical difficulties

Additional tool



Boston Children's Hospital Orthopedic Center









Thank you!





Boston Children's Hospital Orthopedic Center



